

## LA-UR-21-25318

Approved for public release; distribution is unlimited.

Title: Statistical Hauser-Feshbach model calculation for  $(\alpha, n)$  reaction cross section and discrete level population

Author(s): Kawano, Toshihiko

Intended for: technical information exchange with external scientists

Issued: 2021-06-04

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# Statistical Hauser-Feshbach model calculation for $(\alpha, n)$ reaction cross section and discrete level population

Toshihiko Kawano

Theoretical Division  
Los Alamos National Laboratory

June 2, 2021

We calculate the  $\alpha$ -particle induced reactions on  $^{17,18}\text{O}$ ,  $^{19}\text{F}$ , and  $^{23}\text{Na}$ , in the energy range  $0 \leq E_\alpha \leq 10$  MeV, with a particular attention to the branching ratios of populated discrete levels in the  $(\alpha, n)$  channels. Since there are too many open channels to employ the  $R$ -matrix theory, we apply the statistical Hauser-Feshbach model to calculate these reaction cross sections and branching ratios. The branching ratio is defined as

$$b_n = \frac{\sigma_n}{\sum_i \sigma_i + \sigma_c} , \quad (1)$$

where  $\sigma_n$  is the  $n$ -th level production cross section after the neutron emission, and  $\sigma_c$  is the production of the continuum state. In the cases of our target nuclei and the energy range of interest, the residual nuclei of the neutron emission channel are always in their discrete states, so that  $\sigma_c$  can be negligible.

The statistical Hauser-Feshbach code CoH<sub>3</sub> [1, 2] is used to calculate the  $\alpha$ -particle induced reaction. The optical potential parameter for the  $\alpha$ -particle is taken from the work by Avrigeanu et al. [3]. The Koning-Delaroche optical potential [4] is applied to the neutron and proton channels. Because the  $(\alpha, d)$  cross section is too small in the case of this study, we neglected the deuteron channel. Properties of the discrete levels (excitation energy, spin, and parity) are taken from RIPL-3 [5], with some minor modifications based on the most recent ENSDF database [6]. All the other model parameters, such as the level density and the  $\gamma$ -ray strength functions, are the default built-in parameters in CoH<sub>3</sub>. At higher incident energies, a emitted proton sometimes leaves the residual nucleus in its continuum state. Generally speaking, however, the excitation energy is still not so high. The branching ratios are mainly determined by the optical model transmission coefficients and the spin and parity of discrete levels. We also note that the calculated ratios are relatively robust, although the absolute cross sections of discrete level production could have some uncertainties due to the optical potential parameters employed.

The calculated  $(\alpha, n)$  cross sections for  $^{17,18}\text{O}$ ,  $^{19}\text{F}$ , and  $^{23}\text{Na}$  are shown in Fig. 1. Because the statistical model does not predict any resonance structure, these cross sections are understood to be energy-averaged. Besides the  $(\alpha, n)$  cross section, some reaction channels have comparable cross sections to  $(\alpha, n)$ . They are  $^{17}\text{O}(\alpha, \alpha)$  and  $^{19}\text{F}(\alpha, p)$ , which are also shown in this figure.

The calculated branching ratios for  $^{17,18}\text{O}$ ,  $^{19}\text{F}$ , and  $^{23}\text{Na}$  are shown in Fig. 2. Wilson et al. [7] performed the similar calculations with the Hauser-Feshbach code GNASH [8, 9] in the past. There is another model prediction by Lessor and Schenter [10], which is based on some available experimental data. We compare our calculated results with these former predictions in Figs. 3-6.




Figure 1: Calculated  $(\alpha, n)$  reaction cross sections for  $^{17,18}\text{O}$ ,  $^{19}\text{F}$ , and  $^{23}\text{Na}$ , shown by the thick curves. The partial cross sections for the first five levels,  $\sigma_n$ ,  $n = 0-4$ , are shown by the thin curves.




Figure 2: Calculated branching ratios for the  $^{17,18}\text{O}$ ,  $^{19}\text{F}$ , and  $^{23}\text{Na}(\alpha, n)$  reactions to the discrete states in  $^{20,21}\text{Ne}$ ,  $^{22}\text{Na}$ , and  $^{26}\text{Al}$ .




Figure 3: Calculated branching ratios of the  $^{17}\text{O}(\alpha, n)$  reaction to the discrete states in  $^{20}\text{Ne}$ .




Figure 4: Calculated branching ratios of the  $^{18}\text{O}(\alpha, n)$  reaction to the discrete states in  $^{21}\text{Ne}$ .




Figure 5: Calculated branching ratios of the  $^{19}\text{F}(\alpha, n)$  reaction to the discrete states in  $^{22}\text{Na}$ .






Figure 6: Calculated branching ratios of the  $^{23}\text{Na}(\alpha, n)$  reaction to the discrete states in  $^{26}\text{Al}$ .

## References

- [1] T. Kawano, P. Talou, M.B. Chadwick, and T. Watanabe. Monte carlo simulation for particle and  $\gamma$ -ray emissions in statistical hauser-feshbach model. *Journal of Nuclear Science and Technology*, 47(5):462 – 469, May 2010.
- [2] T. Kawano. CoH<sub>3</sub>: The Coupled-Channels and Hauser-Feshbach Code. *Springer Proceedings in Physics*, 254:28 – 34, 2021. CNR2018: International Workshop on Compound Nucleus and Related Topics, LBNL, Berkeley, CA, USA, September 24 – 28, 2018, J. Escher, Y. Alhassid, L.A. Bernstein, D. Brown, C. Fröhlich, P. Talou, W. Younes (Eds.).
- [3] V. Avrigeanu, M. Avrigeanu, and C. Mănăilescu. Further explorations of the  $\alpha$ -particle optical model potential at low energies for the mass range  $A \approx 45$ –209. *Phys. Rev. C*, 90:044612, Oct 2014.
- [4] A. J. Koning and J.-P. Delaroche. Local and global nucleon optical models from 1 keV to 200 MeV. *Nuclear Physics A*, 713(3):231 – 310, 2003.
- [5] R. Capote, M. Herman, P. Obložinský, P. G. Young, S. Goriely, T. Belgya, A. V. Ignatyuk, A. J. Koning, S. Hilaire, V. A. Plujko, M. Avrigeanu, O. Bersillon, M. B. Chadwick, T. Fukahori, Z. Ge, Y. Han, S. Kailas, J. Kopecky, V. M. Maslov, G. Reffo, M. Sin, E. Sh. Soukhovitskii, and P. Talou. RIPL - reference input parameter library for calculation of nuclear reactions and nuclear data evaluations. *Nuclear Data Sheets*, 110(12):3107 – 3214, 2009.
- [6] J.K. Tuli. Evaluated nuclear structure data file. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 369(2):506 – 510, 1996.
- [7] W.B. Wilson, R.T. Perry, E.F. Shores, W.S. Charlton, T.A. Parish, G.P. Estes, T.H. Brown, E.D. Arthur, M. Bozoian, T.R. England, D.G. Madland, and J.E. Stewart. SOURCES 4C: a code for calculating ( $\alpha$ ,n), spontaneous fission, and delayed neutron sources and spectra. Technical Report LAUR-02-1839, Los Alamos National Laboratory, 2002.
- [8] P. G. Young and E. D. Arthur. GNASH, a pre-equilibrium, statistical nuclear-model code for calculation of cross sections and emission spectra. Technical Report LA-6947, Los Alamos National Laboratory, 1977.
- [9] P. G. Young, E. D. Arthur, and M. B. Chadwick. Comprehensive nuclear model calculations: Introduction to the theory and use of the GNASH code. Technical Report LA-12343-MS, Los Alamos National Laboratory, 1992.
- [10] D.L. Lessor and R.E. Schenter. Neutron spectra from ( $\alpha$ ,n) reactions in plutonium compounds calculated from hauser-feshbach reaction theory. Technical Report BNWL-B-109, Battelle Memorial Institute, Pacific Northwest Laboratory, 1971.